

# ***Implications of threshold relationships for projecting fire-regime responses to climate change***

**Adam M. Young<sup>1</sup>, Philip E. Higuera<sup>2</sup>, John Abatzoglou<sup>1</sup>,  
Paul Duffy<sup>3</sup>, and Feng Sheng Hu<sup>4</sup>**

<sup>1</sup>Univ. of Idaho, <sup>2</sup>University of Montana, <sup>3</sup>Neptune and Co., Inc.,

<sup>4</sup>University of Illinois

amyoung@uidaho.edu

**University of Idaho**  
College of Natural Resources

***Alaska Fire Science Consortium***  
***April 25, 2017***


# Outline

- I. Motivation and research questions**
- II. Methods**
- III. Key results and future implications**




# Fire and climate

- ❖ What controlled past variability in fire activity?
- ❖ What do these controls imply about responses to future climate change?




# Fire and climate



*Westerling (2016) Phil Trans B*

***\*Climate is a major driver of fire activity***




*Bowman et al. (2009) Science*




# Fire and climate

Temperature

RCP8.5: 2046-2065




RCP8.5: 2081-2100





Precipitation

2046-2065 - JJA



2081-2100 - JJA





# How can we anticipate future fire activity?


## Three general approaches

Krawchuk and Moritz (2014) *Environmetrics*

### 1. Fire weather index models




### 2. Statistical-correlative models



Moritz et al. (2012) *Ecospheres*

### 3. Fire-process models




Pfeiffer et al. (2013) *Geosci Model Dev*

Flannigan et al. (2013) *For Ecol Mgmt*


# How can we anticipate future fire activity?

## Fire-climate linkages



*\*Future climate will differ compared to the observational record.*

## Projected changes in fire activity





# Testing model transferability

***\* Validate models with independent data source outside the observational record***

- **Paleoecological reconstructions**
  - **Offer independent records of ecological dynamics over past millennia**

## Global Change Biology

Global Change Biology (2012) 18, 1698–1713, doi: 10.1111/j.1365-2486.2011.02635.x

**No-analog climates and shifting realized niches during the late quaternary: implications for 21st-century predictions by species distribution models**

SAMUEL D. VELOZ\*, JOHN W. WILLIAMS\*, JESSICA L. BLOIS\*, FENG HE†, BETTE OTTO-BLIESNER‡ and ZHENGYU LIU†

*Ecology Letters*, (2008) 11: 357–369

doi: 10.1111/j.1461-0248.2007.01150.x

**LETTER**

**Prediction of plant species distributions across six millennia**

*Pearman et al. (2008)*



# Limitations to future projections

Why might projections be wrong?

## 1. Data biases or errors

- Used to construct or inform statistical models
- e.g., GCM projections

## 2. Changing vegetation and ecosystem dynamics




Changing fire-  
climate relationships

# Nature of fire-climate relationships

***\*Fire-climate relationships are nonlinear and contain thresholds***

***\*Small errors or changes may significantly change predictions***





# Key research questions

- (1) How do threshold relationships impact statistical predictions outside the observational range?**
- (2) How sensitive are predictions to modified fire-climate relationships?**
- (3) What are the implications of using threshold relationships to project 21<sup>st</sup>-century changes?**

# Outline

**I. Motivation and research questions**

**II. Methods**

**1. Statistical modeling**

**2. Model-paleodata comparisons**

**III. Key results and future implications**



# Statistical modeling in Alaska



Ecography 40: 606–617, 2017

doi: 10.1111/ecog.02205

© 2016 The Authors. Ecography © 2016 Nordic Society Oikos

Subject Editor: Jessica Blois. Editor-in-Chief: Miguel Araújo. Accepted 30 March 2016


## Climatic thresholds shape northern high-latitude fire regimes and imply vulnerability to future climate change

Adam M. Young, Philip E. Higuera, Paul A. Duffy and Feng Sheng Hu

**Modeled P(fire) at  
30-yr timescales**

**Spatial variation in  
fire activity**

**Presence/absence  
approach**



# Statistical models

## Boosted Regression Trees (BRTs)

- Machine learning algorithm
- Able to fit complex, nonlinear relationships between response and explanatory variables


Fire Data: Alaska Large Fire Database ([fire.ak.blm.gov](http://fire.ak.blm.gov))

Veg. Data: Circumpolar Arctic Veg. Map ([www.geobotany.uaf.edu/cavm](http://www.geobotany.uaf.edu/cavm))


Climate Data: Scenarios Network for Alaska and Arctic Planning ([www.snap.uaf.edu](http://www.snap.uaf.edu))

## Explanatory Variables (1950-2009)


Temp. Warm. Month




Ann. Moisture Avail. (P-PET)



Topography




Veg. Type







# Model performance




***\*Statistical models explain spatial variability in fire presence/absence***

# Historical fire-climate relationships



***\*Non-linear fire-climate relationships reveal thresholds***

***\*Small climatic changes may result in large fire regime changes***



# Outline

**I. Motivation and research questions**

**II. Methods**

1. Statistical modeling

2. Model-paleodata comparisons

**III. Key results and future implications**




# Predicting fire activity for 850-1850 CE

## Use Global Climate Model (GCMs) experiments

### Step 1: Select “best” GCMs


- I. GISS-E2-R
- II. MPI-ESM-P
- III. MRI-CGCM3

### Step 2: Downscale GCM data for 850-1850 CE




### Step 3: Create 30-yr climatologies in AK (per pixel)





# Alaskan paleofire records



# Alaskan paleofire records



CHAR = Charcoal Accumulation Rate




- Wetland tundra
- Shrub tundra
- Graminoid tundra
- Barren tundra
- Boreal forest
- Fire (1950-2009)

**29 fire-history reconstructions in AK**



# Model-paleodata comparisons

## Example



- Evaluating total number of fires for 850-1850 CE
- NOT evaluating predictions over time

# Limitations to future projections

Why might projections be wrong?

## 1. Data biases or errors

- Used to construct or inform statistical models
- e.g., GCM projections

## 2. Changing vegetation and ecosystem dynamics




Changing fire-  
climate relationships

# Limitations to future projections

## Why might projections be wrong?


### 1. Data biases or errors

- Modify value of temperature threshold



### 2. Changing fire-climate relationships

- Modify shape of relationship






# Modifying fire-climate relationships

*\*Evaluate sensitivity of model predictions to slight changes in original relationships*

## Modify threshold values



## Three Modifications

**+0.50 °C**

**+1.00 °C**

**+1.50 °C**


# Modifying fire-climate relationships

*\*Evaluate sensitivity of model predictions to slight changes in original relationships*

Modify shape of relationships

Orig. —

Modified - - -



# Outline

**I. Motivation and research questions**

**II. Methods**


1. Statistical modeling

2. Model-paleodata comparisons



**III. Key results and future implications**



# Q1: How do thresholds impact statistical predictions outside the observational range?



1950-2009 Mean temperature of the warmest month (°C)




*\*Prediction error varies as a function of threshold proximity*

# Why might projections be wrong?


## 1. Data biases or errors

- Modify value of temperature threshold



## 2. Changing fire-climate relationships

- Modify shape of relationship



# Q2: How sensitive are predictions to modified fire-climate relationships?

## Modified threshold values


T1: +0.50 °C

T2: +1.00 °C

T3: +1.50 °C



Prediction error  
(1950-2009)







# Q2: How sensitive are predictions to modified fire-climate relationships?

## Modified relationship shapes



Prediction error  
(1950-2009)



***\*Uncertainty can arise from even small changes in fire-climate relationships***




# Implications for future predictions

# Implications for future predictions

**Q3: How do nonlinear, threshold relationships impact our ability to predict future conditions?**


## Projected changes in fire activity



*Young et al. (2017) Ecography*

# Implications for future predictions

**Q3: How do nonlinear, threshold relationships impact our ability to predict future conditions?**






# Implications for future predictions

MPI-ESM-LR

1971-2000

2010-2039






# Implications for future predictions

GISS-E2-R

1971-2000

2010-2039



Below Thresh.    Near Thresh.    Above Thresh.


# Implications for future predictions

*\* Threshold-driven uncertainty will vary across AK regions in the 21<sup>st</sup> century.*

What are the spatial patterns?

MPI-ESM-LR

2010-2039



Below Thresh. Near Thresh. Above Thresh.




# Implications for future predictions

*\* Tundra and forest tundra dominate areas of highest uncertainty*




2010-2039





# Implications for future predictions

- \* Tundra and forest tundra dominate areas of highest uncertainty*
- \* Regions also most vulnerable to fire-regime shifts*





# Implications for future predictions

2010-2039




*\* Projections accompanied by higher uncertainty*

2070-2099



*\* Projections have less uncertainty as climate moves beyond thresholds*


Relative change in FRP (Future/Historical)



Below Thresh.    Near Thresh.    Above Thresh.

# Caveats and considerations



- ❖ Only consider one explanatory variable (temperature)
- ❖ Do not consider interactions among different driving variables (e.g., temperature and precipitation)
- ❖ Only looked at one modeling tool (i.e., boosted regression trees)





# Conclusions

- ❖ **Uncertainty varies in relation to threshold proximity, and predictions are sensitive to minor modifications**
- ❖ **Threshold-driven uncertainty will vary across AK regions in the 21<sup>st</sup> century.**
- ❖ **Anticipating fire-regime shifts may be accompanied by less threshold-caused uncertainty at the end of century**



# Acknowledgements


Co-authors and committee members: Drs. Philip E. Higuera, John Abatzoglou, Luigi Boschetti, Paul Duffy, and Feng Sheng Hu



Awards ARC-1023477,1023669



Award 14-3-01-7



Award NNX14AK86H



ILLINOIS  
UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN





# Questions?



# Citations

- Barrett, C. M. et al. 2013. Climatic and land cover influences on the spatiotemporal dynamics of Holocene boreal fire regimes. — *Ecology* 94: 389-402.
- Bowman, D. M. J. S. et al. 2009. Fire in the Earth System. — *Science* 324: 481-484.
- Chipman, M. L. et al. 2015. Spatiotemporal patterns of tundra fires: late-Quaternary charcoal records from Alaska. — *Biogeosciences* 12: 4017-4027.
- Flannigan, M. et al. 2013. Global wildland fire season severity in the 21st century. — *Forest Ecol Manag* 294: 54-61.
- Higuera, P. E. et al. 2009. Vegetation mediated the impacts of postglacial climate change on fire regimes in the south-central Brooks Range, Alaska. — *Ecol Monogr* 79: 201-219.
- Higuera, P. E. et al. 2011. Variability of tundra fire regimes in Arctic Alaska: millennial scale patterns and ecological implications. — *Ecol Appl* 21: 3211-3226.
- Hu, F. S. et al. 2015. Arctic tundra fires: natural variability and responses to climate change. — *Front Ecol Environ* 13: 369-377.
- Hu, F. S. et al. 2010. Tundra Burning in Alaska: Linkages to Climatic Change and Sea-Ice Retreat. — *J Geophys Res-Bioge* 115: G04002.
- IPCC, 2013: *The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, 1535 pp
- Kelly, R. et al. 2013. Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. — *Proceedings of the National Academy of Sciences of the United States of America* 110: 13055-13060.
- Krawchuk, M. A. and Moritz, M. A. 2014. Burning issues: statistical analyses of global fire data to inform assessments of environmental change. — *Environmetrics* 25: 472-481.
- Moritz, M. A. et al. 2012. Climate change and disruptions to global fire activity. — *Ecosphere* 3:
- Pearman, P. B. et al. 2008. Prediction of plant species distributions across six millennia. — *Ecol Lett* 11: 357-369.
- Pfeiffer, M. et al. 2013. A model for global biomass burning in preindustrial time: LPJ-LMfire (v1.0). — *Geosci Model Dev* 6: 643-685.
- Veloz, S. D. et al. 2012. No-analog climates and shifting realized niches during the late quaternary: implications for 21st-century predictions by species distribution models. — *Global Change Biol* 18: 1698-1713.
- Westerling, A. L. 2016. Increasing western US forest wildfire activity: sensitivity to changes in the timing of spring. — *Philosophical Transactions of the Royal Society B-Biological Sciences* 371:
- Young, A. M. et al. 2017. Climatic thresholds shape northern high-latitude fire regimes and imply vulnerability to future change. — *Ecography* 40: 606-617.



# Data Sources

## **Fire Data**

Alaska Large Fire Database. available from Alaska Interagency Coordination Center. <http://fire.ak.blm.gov/>.

## **Observational Climate Data (1950-2009)**

Scenarios Network for Alaska and Arctic Planning (SNAP), University of Alaska. 2015. Historical Monthly and Derived Temperature and Precipitation Products - 2 km CRU TS. Retrieved January 2015 from <https://www.snap.uaf.edu/tools/data-downloads>.

## **Vegetation Data and Ecoregions Map**

Homer, C. et al. 2007. Completion of the 2001 National Land Cover Database for the conterminous United States. — Photogramm Eng Rem S 73: 337-341.

Selkowitz, D. J. and Stehman, S. V. 2011. Thematic accuracy of the National Land Cover Database (NLCD) 2001 land cover for Alaska. — Remote Sens Environ 115: 1401-1407.

Nowacki G. et al. 2001. Ecoregions of Alaska. U.S. Geological Survey Open-File Report 02-297 (map).

Walker, D. A. et al. 2005. The circumpolar Arctic vegetation map. — Journal of Vegetation Science 16: 267-282.

## **Topographic Data**

USGS. 1997. Alaska 300m digital elevation model. Anchorage, AK, U.S. Geological Survey EROS Alaska Field Office. <http://agdcftp1.wr.usgs.gov/pub/projects/dem/300m/akdem300m.tar.gz>.

## **Boosted Regression Tree Modeling**

Elith, J. et al. 2008. A working guide to boosted regression trees. — Journal of Animal Ecology 77: 802-813.

Friedman, J. H. 2001. Greedy function approximation: A gradient boosting machine. — Ann Stat 29: 1189-1232.

Ridgeway G with contributions from others. 2015. gbm: Generalized Boosted Regression Models. R package version 2.1.1.

<http://CRAN.R-project.org/package=gbm>.

## **GCM data<sup>†</sup>**

Taylor, K.E., et al. 2012. An Overview of CMIP5 and the experiment design. Bull. Amer. Meteor. Soc. 93: 485-498

Schmidt, G. A. et al. 2011. Climate forcing reconstructions for use in PMIP simulations of the last millennium (v1.0). — Geosci Model Dev 4: 33-45.

Modeling Center	Institute ID	Model Name
NASA Goddard Institute for Space Studies	NASA GISS	GISS-E2-R
Max Planck Institute for Meteorology	MPI-M	MPI-ESM-LR & MPI-ESM-P
Meteorological Research Institute	MRI	MRI-CGCM3

<sup>†</sup> Data retrieved from Earth System Grid Federation:  
<https://esgf.llnl.gov/>